

Supplemental Table S1. List of strains and plasmids used in this study.

Strain, plasmid	Description ^a	Source or reference
Strains:		
<i>E. coli</i>		
Top10	F ⁻ recA1 endA1 hsdR17(rK ⁻ mK ⁺) supE44 thi-1 gyrA relA1	Invitrogen
Rosetta(DE3)	F ⁻ ompT hsdSB(rB ⁻ mB ⁻) gal dcm (DE3) pRARE (Cm ^r)	Novagen
GM2163	F ⁻ ara-14 leuB6 fhuA31 lacY1 tsx78 glnV44 galK2 galT22 mcrA dcm-6 hisG4 rfbD1 rpsL136 dam13::Tn9 xylA5 mtl-1 thi-1 mcrB1 hsdR2	New England Biolabs
XL10-Gold	Tet ^r Δ(mcrA)183 Δ(mcrCB-hsdSMR-mrr)173 endA1 supE44 thi-1 recA1 gyrA96 relA1 lac Hte [F' proAB lacIqZΔM15 Tn10 (Tet ^r) Amy Cm ^r]	Agilent Technologies
<i>Hfx. volcanii</i>		
DS2	Dead sea isolate	(1)
DS70	DS2 cured of plasmid pHV2	(2)
H26	H26 ΔpyrE2	(3)
HM1041	H26 Δsamp1	(4)
HM1042	H26 Δsamp2	(4)
HM1055	H26 Δsamp3	(4)
HM1052	H26 ΔubaA	(4)
NH02	H26 Δsamp1Δsamp2Δsamp3ΔubaA	This study
YW1005	H26 ΔmsrA	This study
YW1006	H26 ΔmsrB	This study
LR01	H26 ΔmsrAΔmsrB	This study
LR02	H26 Δsamp1Δsamp2Δsamp3ΔmsrA	This study
LR03	H26 Δsamp1Δsamp2Δsamp3ΔmsrAΔubaA	This study
XF124	H26 Δsamp1Δsamp2Δsamp3ΔmsrAΔmoaE	This study
Plasmids:		
pJAM202c	Ap ^r ; Nv ^r ; <i>Hfx. volcanii-E. coli</i> shuttle plasmid, empty vector	(5)
pJAM947	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :flag-samp1	(6)
pJAM556	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :flag-samp1 S85R	(7)
pJAM949	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :flag-samp2	(6)
pJAM957	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :ubaA-strepII	(4)
pJAM977	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :flag-samp3	(8)
pJAM3010	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :msrA-strepII	This study
pJAM3011	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :msrB-strepII	This study
pJAM3202	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :flag-samp1 and msrA-strepII	This study
pJAM3203	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :flag-samp2 and msrA-strepII	This study
pJAM3204	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :flag-samp3 and msrA-strepII	This study
pJAM3212	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :flag-samp1 and msrAc _{13S} -strepII	This study
pJAM3213	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :flag-samp2 and msrAc _{13S} -strepII	This study
pJAM3214	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :flag-samp3 and msrAc _{13S} -strepII	This study
pJAM2275	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :flag-samp2 and msrAc _{16S} -strepII	This study
pJAM2276	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :flag-samp2 and msrAc _{48S} -strepII	This study
pJAM2277	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :flag-samp2 and msrAc _{162S} -strepII	This study
pJAM2283	Ap ^r ; Nv ^r ; pJAM202c carries P2 _{rrnA} :flag-samp2 and msrA _{E56A} -strepII	This study
pET24b	Km ^r , IPTG induced expression plasmid	Novagen
pJAM3200	Km ^r , pET24b carries msrA-strepII	This study
pJAM2273	Km ^r , pET24b carries msrAc _{13S} -strepII	This study
pJAM2284	Km ^r , pET24b carries msrA _{E56A} -strepII	This study
pTA131	Ap ^r ; Nv ^r ; pyrE2-based integration vector	(3)
pJAM3012	Ap ^r ; pTA131 carries ΔmsrA with 540-bp 5' and 3' flanking sequences	This study
pJAM3219	Ap ^r ; pTA131 carries ΔmsrB with 500-bp 5' and 3' flanking sequences	This study
pJAM1114	Ap ^r ; pTA131 carries ΔmoaE with 500-bp 5' and 3' flanking sequences	(4)

^aAp^r, ampicillin resistance; Nv^r, novobiocin resistance; Tet^r, tetracycline resistance; Cm^r, chloramphenicol resistance; Km^r, kanamycin resistance. SAMP1, HVO_2619; SAMP2, HVO_0202; SAMP3, HVO_2177; UbaA, HVO_0558; MoaE, HVO_1864; MsrA, HVO_A0230; MsrB, HVO_2234.

References - Supplementary Table S1

1. **Mullakhanbhai MF, Larsen H.** 1975. *Halobacterium volcanii* spec. nov., a Dead Sea halobacterium with a moderate salt requirement. *Arch Microbiol* **104**:207-214.
2. **Wendoloski D, Ferrer C, Dyall-Smith ML.** 2001. A new simvastatin (mevinolin)-resistance marker from *Haloarcula hispanica* and a new *Haloflexax volcanii* strain cured of plasmid pHV2. *Microbiology* **147**:959-964.
3. **Allers T, Ngo HP, Mevarech M, Lloyd RG.** 2004. Development of additional selectable markers for the halophilic archaeon *Haloflexax volcanii* based on the *leuB* and *trpA* genes. *Appl Environ Microbiol* **70**:943-953.
4. **Miranda H, Nembhard N, Su D, Hepowit N, Krause D, Pritz J, Phillips C, Söll D, Maupin-Furlow J.** 2011. E1- and ubiquitin-like proteins provide a direct link between protein conjugation and sulfur transfer in archaea. *Proc Natl Acad Sci U S A* **108**:4417-4422.
5. **Zhou G, Kowalczyk D, Humbard M, Rohatgi S, Maupin-Furlow J.** 2008. Proteasomal components required for cell growth and stress responses in the haloarchaeon *Haloflexax volcanii*. *J Bacteriol* **190**:8096-8105.
6. **Humbard M, Miranda H, Lim J, Krause D, Pritz J, Zhou G, Chen S, Wells L, Maupin-Furlow J.** 2010. Ubiquitin-like small archaeal modifier proteins (SAMPs) in *Haloflexax volcanii*. *Nature* **463**:54-60.
7. **Dantuluri S, Wu Y, Hepowit NL, Chen H, Chen S, Maupin-Furlow JA.** 2016. Proteome targets of ubiquitin-like samp1ylation are associated with sulfur metabolism and oxidative stress in *Haloflexax volcanii*. *Proteomics* **16**:1100-1110.
8. **Miranda HV, Antelmann H, Hepowit N, Chavarria NE, Krause DJ, Pritz JR, Bäsell K, Becher D, Humbard MA, Brocchieri L, Maupin-Furlow JA.** 2014. Archaeal ubiquitin-like SAMP3 is isopeptide-linked to proteins via a UbaA-dependent mechanism. *Mol Cell Proteomics* **13**:220-239.